

Contact heterogeneities in feral swine: implications for disease management and future research

KIM M. PEPIN,^{1,†} AMY J. DAVIS,¹ JAMES BEASLEY,² RAOUL BOUGHTON,³ TYLER CAMPBELL,⁴
SUSAN M. COOPER,⁵ WES GASTON,⁶ STEVE HARTLEY,⁷ JOHN C. KILGO,⁸ SAMANTHA M. WISELY,⁹
CHRISTY WYCKOFF,^{10,11} AND KURT C. VERCAUTEREN¹

¹National Wildlife Research Center, United States Department of Agriculture, 4101 Laporte Avenue, Fort Collins, Colorado 80526 USA

²Savannah River Ecology Laboratory, Warnell School of Forestry and Natural Resources, University of Georgia,
PO Drawer E, Aiken, South Carolina 29802 USA

³Wildlife Ecology and Conservation, Range Cattle Research and Education Center, University of Florida,
3401 Experiment Station, Ona, Florida 33865 USA

⁴East Foundation, 200 Concord Plaza Drive, Suite 410, San Antonio, Texas 78216 USA

⁵Texas A&M AgriLife Research, 1619 Garner Field Road, Uvalde, Texas 78801 USA

⁶USDA/APHIS/Wildlife Services, 602 Duncan Drive, Auburn, Alabama 36849 USA

⁷United States Geological Survey, National Wetlands Research Center, 700 Cajundome Boulevard, Lafayette, Louisiana 70506 USA

⁸Southern Research Station, USDA Forest Service, P.O. Box 700, New Ellenton, South Carolina 29809 USA

⁹Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida 32611 USA

¹⁰Santa Lucia Conservancy, 26700 Rancho San Carlos Road, Carmel, California 93923 USA

¹¹Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, 955 University Boulevard, Kingsville, Texas 78363 USA

Citation: Pepin, K. M., A. J. Davis, J. Beasley, R. Boughton, T. Campbell, S. M. Cooper, W. Gaston, S. Hartley, J. C. Kilgo, S. M. Wisely, C. Wyckoff, and K. C. VerCauteren. 2016. Contact heterogeneities in feral swine: implications for disease management and future research. *Ecosphere* 7(3):e01230. 10.1002/ecs2.1230

Abstract. Contact rates vary widely among individuals in socially structured wildlife populations. Understanding the interplay of factors responsible for this variation is essential for planning effective disease management. Feral swine (*Sus scrofa*) are a socially structured species which pose an increasing threat to livestock and human health, and little is known about contact structure. We analyzed 11 GPS data sets from across the United States to understand the interplay of ecological and demographic factors on variation in co-location rates, a proxy for contact rates. Between-sounder contact rates strongly depended on the distance among home ranges (less contact among sounders separated by >2 km; negligible between sounders separated by >6 km), but other factors causing high clustering between groups of sounders also seemed apparent. Our results provide spatial parameters for targeted management actions, identify data gaps that could lead to improved management and provide insight on experimental design for quantitating contact rates and structure.

Key words: contact; disease transmission; feral swine; GPS; meta-analysis; network; social structure; *Sus scrofa*.

Received 17 July 2015; **accepted** 8 September 2015. Corresponding Editor: A. W. Park.

Copyright: © 2016 Pepin et al. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

† **E-mail:** kim.m.pepin@aphis.usda.gov

INTRODUCTION

Most wildlife populations exhibit contact heterogeneity, especially populations where territorial, dominance or other social grouping behaviors are common. Contact heterogeneity

can alter disease dynamics, changing the timing and peak numbers of infectious individuals (Bansal et al. 2007). The importance of contact heterogeneities in disease transmission has led to a flurry of research to quantitate contact in wildlife populations using a variety of methods

(Cross et al. 2012, Lavelle et al. 2014, Long et al. 2014). Global positioning system (GPS) devices and ultra-high-frequency proximity loggers (PL) are two of the most commonly used technologies for quantitating contact in wildlife. Both devices are used to quantitate contact rates, social networks, factors driving contact heterogeneities and impacts of contact heterogeneities on disease transmission and management (e.g., Cross et al. 2012, Hamede et al. 2012, Drewe et al. 2013, Lavelle et al. 2014, Long et al. 2014, Podgorski et al. 2014, Williams et al. 2014).

Networks are one method for analyzing contact heterogeneities using GPS or PL data from wildlife populations. Network properties that describe contact heterogeneities (e.g., degree, transitivity and connectedness) can be quantitated, and their effects on disease transmission can be studied using disease transmission models (e.g., Bansal et al. 2007, Ames et al. 2011). Degree is the number of individuals to which an individual is connected. Variability in degree distribution affects epidemiological progression by increasing rates of disease spread early during an epidemic and decreasing it later on, relative to a randomly mixing population (Bansal et al. 2007). Networks with some high-degree individuals termed “superspreaders” can reach peak numbers of cases faster and have higher probability of pathogen extinction before an epidemic starts (Lloyd-Smith et al. 2005). Network transitivity describes the proportion of the network that contains individuals whose contacts are interconnected (forming closed triangles: A contacts B and C, and B and C contact each other). Global transitivity can reduce disease transmission because redundancy in local connections leads to local depletion of susceptible individuals (Eames and Keeling 2003). Connectedness describes how completely a network is connected, for example, unconnected networks include one to several discrete clusters with no possibility for disease transmission between them. Realistic representations of contact heterogeneities in wildlife populations can be used to optimize disease management in terms of when and where to increase surveillance and implement controls (Hamede et al. 2012).

Feral swine: data-poor disease threats

Feral swine (*Sus scrofa*) are a highly socially structured species that are globally distributed and pose a threat to human and livestock health

in many countries. They have been implicated in the spillover of economically devastating livestock pathogens such as classical and African swine fever viruses and *Brucella abortus* (Penrith et al. 2011, Costard et al. 2013, Kreizinger et al. 2014), as well as zoonotic pathogens such as hepatitis E virus and influenza A virus (Feng et al. 2014, Caruso et al. 2015). Although the ecological capacity of wild pig populations to maintain pathogens or spark outbreaks is uncertain, population density and social structure are thought to be important due to their effects on contact between individuals (Cowled and Garner 2008, Penrith et al. 2011, Costard et al. 2013).

To our knowledge, only a single study in one location (Poland) has attempted to quantitate contact heterogeneities of feral swine (or wild boar; Podgorski et al. 2014). The study found significant clustering of individuals and sex-based differences in the duration of associations. In addition, it is well-accepted that dominant boars generally occur alone, while reproductively active sows and their offspring occur in groups called sounders (Mayer and Brisbin 2009). Individuals in the same sounder likely contact each other more frequently than individuals from different sounders, but the relative difference in contact rates within and between sounders, and contact structure among sounders is unknown.

To begin addressing these knowledge gaps, we analyzed GPS data from 207 individual pigs (104 males, 103 females) from 11 populations across the United States (Gaston et al. 2008, Campbell and Long 2010, Cooper et al. 2010, Wyckoff et al. 2012, Hartley et al. 2015; and *unpublished data*; Supplement S1: Table S1). All except for one study included individuals from different sounders allowing us to focus on variation in between-sounder contact rates; the scale of contact that is likely most heterogeneous, and potentially limiting disease transmission most strongly. Goals of the above studies were to estimate contact rates with domestic livestock or quantitate pig movement behavior and territoriality. Thus, our analyses of the data should not be taken as absolute measures of contact rates or heterogeneities. Our goals were to: (1) explore potential causes of contact heterogeneities in feral swine; (2) examine the relationship between contact and spatial distribution of home ranges; (3) provide insight

for experimental design of contact studies; and (4) evaluate the effects of missing data on contact structure.

METHODS

We calculated pairwise co-location rates for all pairs of individuals co-monitored for ≥ 7 d, with home range centroids less than 10 km apart. Home range centroids were the median latitude and longitude during co-monitoring. Of the 207 individuals (812 unique pairs), only 3 (three unique pairs) were from the same sounder, thus our analyses focused on contact rates among sounders. Daily co-location rates were calculated as the number of co-locations in a 24-h period between individuals A and B within time interval T and distance D (fixed at 10 m; distance criterion for co-location):

$$\frac{\left(\sum_{j=1}^{n_j} \sum_{k=1}^{n_k} [(d_{j,k} \leq D) \& (t_{j,k} \leq T_x)]\right)}{\text{Number of days overlap}} \quad (1)$$

where $d_{j,k}$ is the distance between location j for individual A and location k for individual B; and $t_{j,k}$ is the time between location j for individual A and location k for individual B (n_j is the total number of locations for individual A and n_k is the total number of locations for individual B). T_x was the time interval between co-locations, where $x = 1$ week or 15 min. These intervals were chosen because they reflect the most frequent time interval that GPS locations were taken (“direct contact” = 15 min between co-locations) and realistic persistence times for bacterial and viral pathogens outside the host (“indirect contact” = 1 week between co-locations). In addition, the mean number of daily locations (location intensity) and duration of monitoring were calculated to account for sampling effort variation. Location intensity was calculated from the mean total locations for sounders A and B throughout the entire period of monitoring (i.e., $n_j + n_k/2$), divided by the number of days they were monitored (i.e., $[n_j + n_k/2]/\text{days of overlap}$).

For statistical analyses we used linear mixed effects models implemented via “lme4” package (R software, R Core Team 2013), using maximum likelihood for parameter estimation. Because we were only interested in factors explaining

variation in contact rates for pairs that made contact at least once, we excluded the pairs that never made contact (i.e., zeros) in all statistical analyses. To account for error correlation within individuals, two random effects were included: unique identifiers for each of collar A (interactor) and collar B (interactee). Our conceptual approach to analyzing fixed effects was hierarchical because we expected that factors intrinsic to the study design (i.e., location intensity and duration of monitoring) would explain significant variation in contact rates, yet we were not interested in their effects. Thus, first we investigated the effects of inherent factors with the goal of removing variation from these factors prior to examining effects of factors we were interested in. Duration of monitoring did not improve AIC over an intercept-only model (Table 1), thus we did not consider this covariate further. In contrast, location intensity significantly improved AIC, thus we used the residuals from the location intensity model as the response in models with covariates such as distance between home range centroids, sex, age class (adult vs. juvenile/subadult) and sounder membership (Table 1). In addition, because distance between home range centroids is another factor that is intrinsic to the study design, we looked at the effects of sex, age and sounder membership both before and after accounting for effects of distance between home range centroids (i.e., response was residuals from a model with just location intensity vs. residuals from a model with location intensity and distance between home range centroids; Table 1). The distance between home range centroids, sex, age class and sounder membership covariates were examined in separate, univariate models because each variable had different amounts of missing data and we were trying to maximize the amount of data used to examine each factor. Although some of the data sets distinguished three age classes (juvenile, subadult and adult), others only distinguished juvenile and adult. Also, there was a lot of missing age-class data. As we were most interested in the age classes that represent pre- and postdispersal, we aggregated the data to make two age classes: adult vs. younger than adult. For models with contact rate data as the response, the response was log-transformed. For all fitted models, the residuals were normally distributed as evidenced by normal quantile plots of the residuals. We calculated

Table 1. Results of statistical analyses. All models included two random effects: unique identifiers for each of interactor A and interactee B. The response and the number of data points differ for Models 5 to 11 thus AIC is not applicable (NA). †

	Response	Fixed effect	Direct contact (within 15 min)				Indirect contact (within 1 week)					
			ΔAIC‡	N	$\beta \pm SE$	<i>P</i>	abs GOF	ΔAIC‡	N	$\beta \pm SE$	<i>P</i>	abs GOF
Test for effects of intrinsic factors												
0	Log contact rate	Int	0	118	-1.31 ± 0.29	<0.0001	0.97	0	193	0.13 ± 0.34	0.69	0.82
1	Log contact rate	LI	-4	118	0.032 ± 0.013	0.013	0.96	-7.9	193	0.05 ± 0.015	0.001	0.80
2	Log contact rate	DM	1.9	118	-0.0014 ± 0.006	0.81	0.97	2	193	-0.002 ± 0.007	0.80	0.82
3	Log contact rate	Dist	-27.5	118	-0.96 ± 0.16	<0.0001	0.97	-35.7	193	-1.43 ± 0.22	<0.0001	0.82
4	Log contact rate	LI + Dist	-33.9	118	0.034 ± 0.011 -0.97 ± 0.15	0.0029 <0.0001	0.97	-56.1	193	0.067 ± 0.013 -1.59 ± 0.21	<0.0001 <0.0001	0.82
Test for effects of ecological variables after accounting for effects of intrinsic factors												
5	Res of 1	Dist	NA	118	-0.17 ± 0.04	<0.0001	0.12	NA	193	-0.73 ± 0.12	<0.0001	
6	Res of 1	Sex	NA	114	-0.19 ± 0.12 -0.22 ± 0.13	0.12 0.088	0.03	NA	181	-0.43 ± 0.35 -0.30 ± 0.38	0.20 0.43	
7	Res of 4	Sex	NA	114	-0.055 ± 0.12 -0.13 ± 0.12	0.64 0.30	0.01	NA	181	-0.15 ± 0.30 -0.06 ± 0.33	0.61 0.86	
8	Res of 1	Age	NA	70	-0.21 ± 0.15 0.049 ± 0.21	0.16 0.82	0.04	NA	117	-0.90 ± 0.43 -0.28 ± 0.52	0.036 0.60	
9	Res of 4	Age	NA	70	-0.17 ± 0.14 0.009 ± 0.20	0.23 0.96	0.02	NA	117	-0.65 ± 0.37 -0.097 ± 0.44	0.076 0.83	
10	Res of 1	Mem	NA	11	0.65 ± 0.16	<0.0001	0.94	NA	20	1.83 ± 0.68	0.007	
11	Res of 4	Mem	NA	11	0.93 ± 0.08	<0.0001	0.98	NA	20	1.28 ± 0.42	0.0023	

† Abbreviations are: Res, residuals; LI, location intensity; DM, duration of monitoring; Dist, distance between home range centroids; Age, age class with three factors (immature–immature, immature–adult, adult–adult); Sex, includes three factors (female–female, male–female, male–male); Mem, sounder membership (indicator for belonging to the same sounder); SE, standard error of the estimate; abs GOF, absolute goodness of fit; squared correlation coefficient between observed and predicted values.

‡ ΔAIC's are relative to the null model with no fixed effects; negative values of >2 represent a significant improvement over the null model.

P-values using the *t* statistics generated by the “lmer” function (lme4 package, R Software), assuming that the *t*-distribution converges to the *z* distribution at large sample size. We assessed explained variation in the models (absolute goodness of fit) with squared correlation coefficients of observed and model-predicted data.

We calculated network properties (mean degree, global transitivity and number of independent clusters) using the “igraph” package in R (Csardi and Nepusz 2006). Network properties were based on undirected, unweighted graphs (i.e., binary adjacency matrices), except we additionally calculated transitivity from weighted graphs to examine how contact rate heterogeneity affected this property. Mean degree was the average number of edges per node. Global transitivity for the unweighted networks was the ratio of triangles to

triplets in the network (Csardi and Nepusz 2006). A triangle was defined as a closed triplet (three nodes connected by three edges), whereas a triplet was defined as three nodes connected by either two or three edges (thus a triangle is a special case of a triplet). Global transitivity for the weighted networks was calculated using the “clustering_w” function in the “tnet” package in R, and assuming the geometric mean method for averaging weights, as defined in Opsahl and Panzarasa (2009). The equation for calculating global transitivity of the weighted networks is equivalent to the ratio of triangles to triplets when an unweighted network is used. The number of independent clusters was the total number of either single or subgroups of connected nodes that were completely unconnected to the rest of the network. The number of independent clusters was calculated using the “clusters”

function in the “igraph” package. For the network analyses, we only included three studies with the most individuals ($N = 20, 18$ and 13). We also dropped different amounts of data (from 1 point up to 50% of points) from the networks and recalculated the network properties to investigate effects of missing data. For each number of data points excluded, nodes were dropped at random and the procedure was repeated 100 times.

RESULTS AND DISCUSSION

Heterogeneity in co-location rates

The maximum daily contact rates among sounders were 45 times higher for indirect relative to direct contact. Direct contact rates did not change with the duration of monitoring (Fig. 1a, c), but did increase significantly with the number of locations recorded per day ($P = 0.013$ for direct contact and $P = 0.001$ for indirect contact; Fig. 1b, d, Models 1 and 2 in Table 1). This emphasizes that when using GPS-type data for quantitating contact rates, the interval between locations should approximate the minimum time between the start and end of a perceived disease-relevant contact. For example, if a pathogen requires 5 min of contact for a high probability of transmission, then to quantitate disease-relevant contact rates recording locations at least every 5 min would be important. Furthermore, variability from the duration of monitoring may be weak or insignificant except when durations are very short (<7 d in our case). Additional studies, which include long-term monitoring should be undertaken to determine if contact rates change seasonally, which could obscure the potential importance of monitoring duration on quantitation of contact rates.

Effects of distance between home range centroids on contact rates

As previous work has shown that wild pig home ranges are relatively small, we expected that distance between pig home range centroids would be a strong predictor of contact rates. Indeed, contact rates decreased significantly with distance between home range centroids (Fig. 2a; considering pairs with at least one contact only: $P < 0.0001$ for direct and indirect contact, Model 3 in Table 1), and this relationship remained significant even after variation from location intensity was removed ($P < 0.0001$

for direct and indirect contact, Model 5 in Table 1). Contacts were rare between individuals in different sounders whose home range centroids were >2 km, despite there being many data points from pairs of sounders with home range centroids separated by further distances (up to ~ 10 km; Fig. 2a). There were very few contacts for pairs with home range centroids separated by 2–6 km and none for pairs at distances >6 km. Thus, direct disease transmission is likely to be rare between sounders ranging further than 2 km and non-existent between sounders separated by 6 km, unless a significant amount of long-distance dispersal occurs during persistence of a given disease.

Effects of sex and age class on contact rates

No significant effects of sex on either direct (Fig. 2b) or indirect contact were apparent (Models 6 and 7, Table 1) after accounting for intrinsic factors, suggesting that any potential sex-based differences may be mostly explained by intrinsic factors. Effects of age class were significant on indirect contact but not direct contact (Fig. 2c), after variation from location intensity was accounted for (Model 8 in Table 1), but not after variation from both location intensity and distance was considered (Model 9 in Table 1). Specifically, indirect contacts between different age classes were significantly lower than indirect contacts between like age classes. The age class results suggest that although age-based differences in contact rates may explain some variation in indirect contact rates among sounders, it may be less significant for explaining variation in direct contact rates. Also, the variation in contact rates among sounders that was explained by age class was very small (explained variation = 0.04) and when distance between home range centroids was accounted for, the age class effect became insignificant (variation explained was reduced by 50%), suggesting that potential heterogeneities due to age class could be mostly explained by other factors such as distance between home range centroids. Although it should be noted that age class data were missing in some studies and ages were defined using different methods across studies, emphasizing that effects of age class should be tested more rigorously in a study that uses a standardized measure of age for all individuals




Fig. 1. Co-locations for each dyad in all studies. Only pairs that had at least one contact are shown. Points are coded by different markers and colors to distinguish studies in all plots. (a, c) Log daily contact rates plotted against the number of days that pairs were monitored. (b, d) Log daily contact rates plotted against the daily frequency of recording locations (location intensity). (a, b) direct contact; and (c, d) indirect contact.

and designed to decipher the relative contributions of age-based differences in contact rates compared with home range locations.

Effects of sounder membership on contact rates

Although only three individuals in one study (SC) were from the same sounder, sounder membership explained a significant amount of variation in both direct (Fig. 2d) and indirect contact rates even after location intensity and distance between home range centroids were accounted for (Models 10 and 11 in Table 1). Fig. 2d shows a >20-fold difference in contact rates for pairs from the same vs. adjacent sounders. This difference was roughly fivefold for indirect contact.

Taken together, our results emphasize that heterogeneities in contact rates are substantial in feral swine populations. Most of the variation can be explained by individual, location intensity, dis-

tance between home range centroids and sounder membership. Although sex was not significant and age class only explained a tiny amount of variation, it may be worth testing effects of these factors in future studies that are specifically designed to measure contact rates. The random effects for interactor A and interactor B explained the largest proportion of variation (compare absGOF for Model 0 to Models 1–4), highlighting that some individuals are inherently more likely to interact with others. After accounting for location intensity, distance between home range centroids and sounder membership were the two most important factors explaining variation in contact rates after individual-level behavior.

Contact structure

In all three of the studies for which we calculated network properties (FL, TX4 and TX6),




Fig. 2. Factors affecting co-location rates between sounders. Direct contact rates. X-axes are: distance between home range centroids in km (a, d) sex of the individuals in a pair (b: FF, female–female; MF, male–female; MM, male–male), age class of the individuals in a pair (c: JJ, juvenile–juvenile; JA, juvenile–adult; AA, adult–adult; note – “juvenile” includes subadults). Points are color-coded by study and scaled to the numbers of days that the pair was monitored. (a, d) Size of points is scaled to the location intensity. (b, c) Size of points is correlated with the number of pairs. Numbers along the top show the total number of pairs in each category. Error bars are the standard deviations of the mean contact rates on a linear scale.

average degree (G) and connectedness (C) increases (meaning the value of C decreases) as the time between contacts increases (meaning contact is more indirect, Fig. 3), highlighting that consideration of indirect contacts increases opportunity for disease transmission through increased connectedness between individuals (or, here, sounders; Drewe et al. 2013). Network properties differed across studies and contact type (direct vs. indirect) despite the relatively stable pattern in average distance between home range centroids (d , Fig. 3), suggesting that network structure may describe contact heterogeneities in addition to those imposed by spatial relationships.

Global transitivity for the unweighted networks decreased with increasing time between

contacts in two of the studies (FL and TX4), but it showed an inconsistent pattern in the third study (TX6, see T values Fig. 3). This inconsistency may partly be due to the very weak connectedness of these highly disconnected networks (up to nine distinct clusters), i.e., transitivity levels likely vary between distinct clusters and thus global effects from addition of connections strongly depend on which connections are added. It is also noteworthy that transitivity was very high (≥ 0.67 for all networks) and connectedness very low in all three networks from geographically distinct populations. When transitivity was calculated by considering contact rate heterogeneities, it was even higher at ≥ 0.95 for all networks, suggesting that high transitivity may be a common characteristic in wild pig populations. This hypothesis




Fig. 3. Spatial networks for three studies: FL (green), TX4 (red), TX6 (black). All networks were scaled to a 10×10 km grid; scaling is shown in the lower right plot. Networks are plotted for three different types of contact: direct (within 15 min), indirect (within 1 week), and indirect (within 1 month). Data in the plots are: d = mean distance (± 1 SD) between connected nodes, G = degree (± 1 SD), R_u = global transitivity for unweighted networks, R_w = global transitivity for weighted networks, C = number of independent clusters.

should be validated with a study that is specifically designed to measure contact networks because disease dynamics in disconnected, clustered populations are predicted to be very different from those in fully connected, unclustered ones and disease management is most effective when based on knowledge of contact structure (Eames and Keeling 2003, Hamede et al. 2012). One biological explanation for the high transitivity could be that sounders are more dynamic, frequently swapping members between sounders with closely related individuals, than is traditionally believed. To verify this hypothesis, all individuals from multiple adjacent sounders need to be monitored for contact, a study that remains to be done.

The networks we presented are based on a small number of sounders (13–20), undoubtedly

with missing data (i.e., not all sounders within the area of the sampled individuals were included). We explored the effects of such missing data on network properties by dropping up to 50% of the sounders and recalculating the network properties (Fig. 4). Degree decreased significantly when 50% of nodes were dropped (Fig. 4). Transitivity was relatively stable with up to 50% missing data for the networks with indirect contacts, but changed in different directions in the direct-contact networks (which were much more clustered). The number of independent clusters was also more sensitive to missing data in the direct-contact networks relative to those with indirect contacts. Thus, networks of wild pig social structure based on indirect contacts may be relatively robust to inherent difficulties with field




Fig. 4. Effects of missing data on network properties. The three studies with the most connections are shown. Points are scaled by the number of nodes (FL = 18, TX4 = 13, TX6 = 20). Rows: mean connections per node, global transitivity (based on unweighted network), and number of independent clusters. Columns are types of contact: direct (within 15 min), indirect (within 1 week), and indirect (within 1 month). Error bars are 2 SDs of 100 replicates. Stars show values where 50% of nodes are dropped: 9 – FL, 6.5 – TX4, 10 – TX6).

study design. Nevertheless, to quantitate contact structure every effort should be made to sample all adjacent sounders within the target spatial area.

CONCLUSION

The 45-fold variation in direct vs. indirect contact rates, and differences in network properties, highlight that appropriately defining disease-specific direct and indirect contact is a crucial first step for designing studies to measure contact rates. To quantitate contact rates in absolute terms for predicting rates of disease spread, a second important consideration is to record locations at a frequency that approximates the minimum time of an effective contact. As expected, within-sounder contact rates appear to be much higher than between-sounder rates (>20 times in our study), but studies with larger sample sizes are needed to quantitate the magnitude of

these differences. Between-sounder contact rates strongly depended on home range location, suggesting that between-sounder contact heterogeneities can be partly described using knowledge of wild pig movement ecology. Specifically, we expect disease transmission to be reduced between sounders separated by >2 km and negligible between sounders separated by >6 km. It should be noted, however, that densities of the populations sampled are unknown, and lower density populations could show increased contact across greater distances—a relationship that remains to be studied. In addition to space, other factors causing high clustering may be at play (e.g., access to water, baiting). Studies designed to identify and quantitate these factors, as well as potential seasonality, are imperative for planning effective management of disease threats from feral swine because they directly impact the timing, severity and spatial spread of outbreaks.

ACKNOWLEDGMENTS

We thank two anonymous reviewers for their careful, insightful reviews of an earlier version of this manuscript. Funding was provided by the U.S. Department of Agriculture National Wildlife Research Center and the U.S. Department of Energy under Award Number DE-FC09-07SR22506 to the University of Georgia Research Foundation and under Interagency Agreement No. DE-AI09-00SR22188 with the U.S. Department of Agriculture Forest Service Savannah River. USDA NWRC provided funding to University of Florida under USDA APHIS Cooperative Agreement 1274120896CA.

LITERATURE CITED

- Ames, G. M., D. B. George, C. P. Hampson, A. R. Kanarek, C. D. McBee, D. R. Lockwood, J. D. Achter, and C. T. Webb. 2011. Using network properties to predict disease dynamics on human contact networks. *Proceedings of the Royal Society B-Biological Sciences* 278:3544–3550.
- Bansal, S., B. T. Grenfell, and L. A. Meyers. 2007. When individual behaviour matters: homogeneous and network models in epidemiology. *Journal of the Royal Society, Interface* 4:879–891.
- Campbell, T. A., and D. B. Long. 2010. Activity patterns of wild boars (*Sus scrofa*) in southern Texas. *Southwestern Naturalist* 55:564–567.
- Caruso, C., et al. 2015. Serological and virological survey of hepatitis E virus in wild boar populations in northwestern Italy: detection of HEV subtypes 3e and 3f. *Archives of Virology* 160:153–160.
- Cooper, S. M., H. M. Scott, G. R. de la Garza, A. L. Deck, and J. C. Cathey. 2010. Distribution and interspecies contact of feral swine and cattle on rangeland in south Texas: implications for disease transmission. *Journal of Wildlife Diseases* 46:152–164.
- Costard, S., L. Mur, J. Lubroth, J. M. Sanchez-Vizcaino, and D. U. Pfeiffer. 2013. Epidemiology of African swine fever virus. *Virus Research* 173:191–197.
- Cowled, B., and G. Garner. 2008. A review of geospatial and ecological factors affecting disease spread in wild pigs: considerations for models of foot-and-mouth disease spread. *Preventive Veterinary Medicine* 87:197–212.
- Cross, P. C., T. G. Creech, M. R. Ebinger, D. M. Heisey, K. M. Irvine, and S. Creel. 2012. Wildlife contact analysis: emerging methods, questions, and challenges. *Behavioral Ecology and Sociobiology* 66:1437–1447.
- Csardi, G., and T. Nepusz. 2006. The igraph software package for complex network research. *Inter Journal, Complex Systems* 1695:1–9; <http://igraph.org>.
- Drewe, J. A., H. M. O'Connor, N. Weber, R. A. McDonald, and R. J. Delahay. 2013. Patterns of direct and indirect contact between cattle and badgers naturally infected with tuberculosis. *Epidemiology and Infection* 141:1467–1475.
- Eames, K. T. D., and M. J. Keeling. 2003. Contact tracing and disease control. *Proceedings of the Royal Society B-Biological Sciences* 270:2565–2571.
- Feng, Z. X., et al. 2014. Influenza A Subtype H3 viruses in feral swine, United States, 2011–2012. *Emerging Infectious Diseases* 20:843–846.
- Gaston, W., J. B. Armstrong, W. Arjo, and H. L. Stribling. 2008. Home range and habitat use of feral hogs (*Sus scrofa*) on Lowndes County WMA, Alabama. National Conference on Feral Hogs, St. Louis, Missouri.
- Hamede, R., J. Bashford, M. Jones, and H. McCallum. 2012. Simulating devil facial tumour disease outbreaks across empirically derived contact networks. *Journal of Applied Ecology* 49:447–456.
- Hartley, S. B., B. L. Goatcher, and S. K. Sapkota. 2015. Movements of wild pigs in Louisiana and Mississippi, 2011–13. Page 11 *in* US Geological Survey, US Department of the Interior, editors. U.S. Geological Survey Open-File Report 2014–1241. <http://dx.doi.org/10.3133/ofr20141241>.
- Kreizinger, Z., J. T. Foster, Z. Ronai, K. M. Sulyok, E. Wehmann, S. Janosi, and M. Gyuranecz. 2014. Genetic relatedness of *Brucella suis* biovar 2 isolates from hares, wild boars and domestic pigs. *Veterinary Microbiology* 172:492–498.
- Lavelle, M. J., J. W. Fischer, G. E. Phillips, A. M. Hildreth, T. A. Campbell, D. G. Hewitt, S. E. Hygnstrom, and K. C. Vercauteren. 2014. Assessing risk of disease transmission: direct implications for an indirect science. *BioScience* 64:524–530.
- Lloyd-Smith, J. O., S. J. Schreiber, P. E. Kopp, and W. M. Getz. 2005. Superspreading and the effect of individual variation on disease emergence. *Nature* 438:355–359.
- Long, J. A., T. A. Nelson, S. L. Webb, and K. L. Gee. 2014. A critical examination of indices of dynamic interaction for wildlife telemetry studies. *Journal of Animal Ecology* 83:1216–1233.
- Mayer, J. J., and I. L. Brisbin. 2009. Wild pigs: biology, damage, control techniques and management. Department of Energy, editor. http://www.sdmmp.com/Libraries/Management_Plans_and_Reports/SRNL-2009_Wild_Pigs_biology_damage_mgmt.sflb.ashx.
- Opsahl, T., and P. Panzarasa. 2009. Clustering in weighted networks. *Social Networks* 31:155–163.
- Penrith, M. L., W. Vosloo, and C. Mather. 2011. Classical swine fever (hog cholera): review of aspects relevant to control. *Transboundary and Emerging Diseases* 58:187–196.

- Podgorski, T., D. Lusseau, M. Scandura, L. Sonnichsen, and B. Jedrzejewska. 2014. Long-lasting, kin-directed female interactions in a spatially structured wild boar social network. *PLoS ONE* 9:e99875.
- R Core Team. 2013. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>
- Williams, D. M., A. C. Dechen Quinn, and W. F. Porter. 2014. Informing disease models with temporal and spatial contact structure among GPS-collared individuals in wild populations. *PLoS ONE* 9:e84368.
- Wyckoff, A. C., S. E. Henke, T. A. Campbell, D. G. Hewitt, and K. C. VerCauteren. 2012. Movement and habitat use of feral swine near domestic swine facilities. *Wildlife Society Bulletin* 36:130–138.

SUPPORTING INFORMATION

Additional Supporting Information may be found online at: <http://onlinelibrary.wiley.com/doi/10.1002/ecs2.1230/supinfo>